



Contents lists available at SciVerse ScienceDirect

Molecular Phylogenetics and Evolution

journal homepage: www.elsevier.com/locate/ympev

Genetic aspects of mitochondrial genome evolution

Matthias Bernt^a, Anke Braband^b, Bernd Schierwater^{c,d,e}, Peter F. Stadler^{f,g,h,i,j,*}^aParallel Computing and Complex Systems Group, Department of Computer Science, University of Leipzig, Augustusplatz 10, D-04109 Leipzig, Germany^bLGC Genomics GmbH, Ostendstr. 25, 12459 Berlin, Germany^cITZ, Ecology & Evolution, TiHo Hannover, Buenteweg 17d, 30559 Hannover, Germany^dDepartment of Invertebrates, American Museum of Natural History, 79 St. at Central Park West, New York, NY 10024, USA^eDepartment of Molecular, Cellular and Developmental Biology, Yale University, 165 Prospect St., New Haven, CT 06511, USA^fBioinformatics Group, Department of Computer Science, and Interdisciplinary Center for Bioinformatics, University of Leipzig, Härtelstraße 16-18, D-04107 Leipzig, Germany^gMax-Planck-Institute for Mathematics in the Sciences, Inselstraße 22, D-04103 Leipzig, Germany^hFraunhofer Institut für Zelltherapie und Immunologie – IZI, Perlickstraße 1, D-04103 Leipzig, GermanyⁱDepartment of Theoretical Chemistry, University of Vienna, Währingerstraße 17, A-1090 Wien, Austria^jSanta Fe Institute, 1399 Hyde Park Rd., Santa Fe, NM 87501, USA

ARTICLE INFO

Article history:

Available online xxx

Keywords:

Genetic code
Gene content
Chromosome structure
Replication
Transcriptome
Phylogenomics

ABSTRACT

Many years of extensive studies of metazoan mitochondrial genomes have established differences in gene arrangements and genetic codes as valuable phylogenetic markers. Understanding the underlying mechanisms of replication, transcription and the role of the control regions which cause e.g. different gene orders is important to assess the phylogenetic signal of such events. This review summarises and discusses, for the Metazoa, the general aspects of mitochondrial transcription and replication with respect to control regions as well as several proposed models of gene rearrangements. As whole genome sequencing projects accumulate, more and more observations about mitochondrial gene transfer to the nucleus are reported. Thus occurrence and phylogenetic aspects concerning nuclear mitochondrial-like sequences (NUMTS) is another aspect of this review.

© 2012 Elsevier Inc. All rights reserved.

1. Introduction

Mitochondria are organelles with their own genetic material that are present in nearly all eukaryotic cells. In addition to the production of most of a cell's supply of ATP they are involved in many other key processes, such as the control of the cell cycle and cell growth (McBride et al., 2006). They share many features with prokaryotes and are commonly thought to have originated as endosymbionts in the ancestral eukaryote. A recent phylogenomic study suggests a common origin of mitochondria and the SAR11 clade of *Alphaproteobacteria* as a sister group to the *Rickettsiales* (Thrash et al., 2011).

Compared to the mitochondria found in other eukaryotic kingdoms, metazoan mitochondria are massively reduced in their genetic structure. A typical metazoan mitogenome consists of a single circular DNA that carries 13 intron-less protein coding genes, the two subunits of the mitochondrial ribosomal RNA, and 22 tRNAs, one for each amino acid except leucine and serine, which

have two copies each. Almost their complete proteome, much of which is of prokaryotic origin, is imported from the nucleus (Chacinska et al., 2009; Huynen, 2010). The expression of the few proteins encoded on the mitochondrial DNA (mtDNA) uses its own translation system and often employs lineage-specific variations of the genetic code. The mtDNA is not organised in nucleosomal bodies but instead associated with the mitochondrial inner membrane and several proteins including mitochondrial transcription factor A (mtTFA) and mitochondrial single-stranded DNA binding protein (mtSSB) (Bogenhagen et al., 2003). The inheritance of mtDNA is strictly maternal, with the single known exception of the doubly uniparental inheritance in unionid mussels (Breton et al., 2010).

In metazoan animals, the analysis of mitochondrial genomes with respect to size, genome architecture, gene content/arrangement, exon/intron structure, genetic code and secondary structure of rRNA and tRNA genes has been a powerful tool to infer phylogenetic relationships at different taxonomic levels (Boore, 2006). Furthermore, mitochondrial protein coding sequence data harbour valuable phylogenetic signal to further clarify uncertain relationships within and between phyla (Bourlat et al., 2008; Zardoya and Meyer, 1996).

This review is organised as follows: We first provide an overview of the general properties of mitochondrial genomes with a

* Corresponding author at: Bioinformatics Group, Department of Computer Science, and Interdisciplinary Center for Bioinformatics, University of Leipzig, Härtelstraße 16-18, D-04107 Leipzig, Germany. Fax: +49 341 97 16679.

E-mail addresses: bernt@informatik.uni-leipzig.de (M. Bernt), phylogenetics@arcor.de (A. Braband), bernd.schierwater@ecolevol.de (B. Schierwater), studla@bioinf.uni-leipzig.de (P.F. Stadler).

focus on the aberrant structures and the variations of the genetic code in metazoan phyla. In addition we address nuclear copies of mitochondrial DNA (NUMTS). Section 3 is concerned with topics related to replication of mitochondrial genomes, such as strand biases, a review of the models of replication itself, mutation and repair, and mechanisms of genome rearrangements. Transcription of mitogenomes is treated in the last part of the paper.

2. Genomics

2.1. Genome organisation

A typical metazoan mitogenome encodes 13 proteins that belong to four enzyme complexes of the respiration chain: cytochrome b (*cob*), two subunits of ATP synthase (*atp6,8*), three subunits of cytochrome c oxidase (*cox1–3*), and seven subunits of NADH dehydrogenase (*nad1–6, 4L*), see Fig. 1. The mitogenome furthermore contains the small and large (*rrnS, rrnL*) RNA subunits of the mitochondrial ribosomes. These rRNAs are clearly of prokaryotic origin. A comprehensive review of gene content and genome organisation of animal mitochondria can be found in Gissi et al., 2008.

Typically, 22 tRNAs are encoded on the mtDNA. With the exception of serine and leucine, there is only a single tRNA for each amino acid. In many cases one or more mitochondrial tRNAs are missing and functionally replaced by nuclear tRNAs (Alfonzo and Söll, 2009). A peculiar feature are overlapping tRNAs, as in the case of *tRNA-Trp* and *tRNA-Cys* in Arthropoda (Satta et al., 1987). Mitochondrial tRNAs, in contrast to tRNAs in other genetic systems,

do not necessarily fold into the famous clover-leaf structure, although more than 90% of the known cases conform to the standard. Bizarre structures have evolved independently in many metazoan clades. The most frequent aberration is the lack of conserved D-arms (DHU-arm) and/or T-arms (TΨC-arm). In most Bilateria, *tRNA-Ser* (AGN) lacks the D-arm. The corresponding deletion in *tRNA-Ser* (UCN) is prevalent among protostomes. In diverse taxa scattered throughout the Bilateria (e.g. Mammalia, Amphibia, and many Ecdysozoa) furthermore, one finds *tRNAs-Cys* without D-arm. A well known example for a systematic reduction of mt-tRNAs are Nematoda (Wolstenholme et al., 1994). With the exception of the two serine tRNAs, here the T-arm is deleted in all other tRNAs. Even the deletion of both the T- and the D-arm are not uncommon in particular in Enoplea (Jühling et al., 2012b). In Onychophora elaborate editing is capable of repairing large tRNA fragments (Segovia et al., 2011). Aberrant tRNAs structures have been collected and studied systematically as part of a recent effort to re-annotate mitochondrial tRNAs (Jühling et al., 2012a). Like their nuclear counterparts, mt-tRNAs are chemically modified. In some cases the modifications are necessary to ensure correct folding (Helm, 2006). A 5-taurinomethyluridine found in *Halocynthia roretzi* at the anticodon wobble positions of *tRNA-Met* (AUR), *tRNA-Trp* (UGR), and *tRNA-Gly* (AGR) appears to be responsible for the deciphering of the non-universal codes in ascidians.

Only a small fraction of most metazoan mitogenomes does not code for one of the proteins, rRNAs, or tRNAs. In contrast, the mitochondrial genomes of most other Eukarya are much larger and significantly differ in their gene content. While the gene content is extremely well conserved in animals, there are major differences among Metazoa in the arrangements of the few elements that




Fig. 1. Representation of the human mitogenome which has the gene complement typical for metazoan mitogenomes; image generated with *mtviz* <http://pacosy.informatik.uni-leipzig.de/mtviz/>. The coding strand is indicated by thick line; abbreviations are as in the text and for tRNAs the one letter code of the corresponding amino acid is given; for the leucine and serine tRNAs the recognised codon is indicated. The inset sketches the organisation of the control region. The three conserved sequence blocks (CSBs) are marked I, II, and III. The primer for H-strand replication and the two H-strand transcripts are indicated.

regulate replication and transcription. In chordates, the most prominent non-coding region is the “displacement loop” (D-loop). It harbours almost all of the regulatory sequences for replication and transcription, see the inset in Fig. 1. A second non-coding region of only ~30nt located inside a tRNA cluster harbours the origin of replication for the L-strand in vertebrates (Fernández-Silva et al., 2003). Arthropods, and possibly also nematodes, have a single large AT-rich non-coding region which contains the replication origins for both strands (Zhang and Hewitt, 1997; Saito et al., 2005). In many of the major metazoan clades little or nothing is known about the mitochondrial control structures. Mollusca may serve as an example. In Pulmonata, several small non-coding regions with poor conservation have been reported (Yamazaki et al., 1997), while Cephalopoda have large “unassigned regions” that in part contain repetitive sequences (Boore, 2006). Gastropod mitogenomes, on the other hand have several usually small non-coding regions that are AT-rich and contain clover-leaf like structures. One or more of them have been suggested as replication origins (Grande et al., 2008, 2009).

2.2. Aberrant genome structures

Deviations from the typical organisation are relatively rare and appear to be restricted to individual clades. In most cases, there are missing or additional genes, most frequently a loss of tRNA genes. In diploblasts, the deletion of many or even most tRNAs is a common phenomenon (Haen et al., 2010; Wang and Lavrov, 2008). Among protostomes, in particular Rotifera (Suga et al., 2008) and Onychophora (Braband et al., 2010) are affected. Marsupials are well known to have lost mt-*tRNA-Lys* (Dörner et al., 2001). For a recent comprehensive overview of tRNA losses we refer to (Jühling et al., 2012a), see also (Gissi et al., 2008). The protein coding gene that is missing most often is *atp8*. It is absent e.g. in many Nematoda and some Mollusca (Boore, 1999). According to both published annotations, surveyed e.g. in (Gissi et al., 2008), and the re-annotation of the animal genomes with MITOS (Bernt et al., 2012b, in this special issue), *atp8* is lost in scattered cases throughout all major metazoan clades. These data show that the loss in Nematoda is not as wide-spread as previously thought and high-light Platyhelminthes as another hot spot for the loss of *atp8*. Due to the short and rapidly evolving sequence of *atp8* it remains uncertain, however, which of these cases are failures to detect the gene and which are true gene losses. In octocoral mitogenomes an additional *msh1* gene is found (McFadden et al., 2006, and references therein). It has recently been identified as a compelling candidate for horizontal gene transfer and appears to have a function in DNA repair (Bilewicz and Degnan, 2011).

In some cases, however, the genome structures are dramatically different, see Table 1. The mitogenome of the human body louse *Pediculus humanus*, for instance has disintegrated into 18 mini chromosomes with a size of 3–4 kb each containing 1–3 genes. Other sucking lice (but neither chewing lice nor Psocoptera) have a similar genome structure (Shao et al., 2009; Cameron et al., 2011). There is unequivocal evidence for recombination between different mini chromosomes, although the mechanism remains unknown (Shao and Barker, 2011). Multipartite mitogenomes have also been reported for the isopod *Armadillidium vulgare* (Raimond et al., 1999), the rotifer *Brachionus plicatilis* (Suga et al., 2008) and the nematode *Globodera pallida* (Armstrong et al., 2000). In the latter, the gene content between chromosomes is partially redundant and contains repetitive sequences that have been speculated to play a role in replication (Armstrong et al., 2000). Medusozoa (Cnidaria excluding Anthozoa) feature linear chromosomes that are split into two or more different molecules (Voigt et al., 2008; Kayal et al., 2012).

Table 1

Examples of aberrant structures of metazoan mitogenomes.

Taxon	Description	Reference
Phthiraptera	Mini circles	Cameron et al. (2011)
Globodera	Multiple chromosomes	Armstrong et al. (2000)
Armadillidium	Linear + circular	Raimond et al. (1999)
Brachionus	2 Chromosomes	Suga et al. (2008)
Medusozoa	Linear fragments	Kayal et al. (2012)
Porifera	Hairpin-forming repeats	Erpenbeck et al. (2009)
Onychophora	tRNA loss	Podsiadlowski et al. (2008)
Chaetognatha	tRNA loss	Helfenbein et al. (2004)
Cnidaria	tRNA loss	Kayal and Lavrov (2008)
Unionida	F ORFs	Breton et al. (2010)
Placozoa	Several extra ORFs	Dellaporta et al. (2006)
	Group I intron, split genes	Burger et al. (2009)
Porifera	Group I intron	Rot et al. (2006)
Cnidaria	Group I intron	Beagley et al. (1996)
Crassostrea	Fragmented rRNAs	Milbury et al. (2010)
Annelida	Group II intron	Vallès et al. (2008)

At least in mammals, different topological organisations of the mitogenomic DNA have been reported. Besides the usual, circular DNA also linear molecules and a variety of concatenated forms are present in varying amounts (see Pohjoismäki and Goffart, 2011, and references therein). Data for a comparative analysis of phenomenon within Metazoa seem to be absent.

Unionid mussels form an exception from the otherwise strict maternal inheritance of mitochondria. The doubly uniparental inheritance (DUI) found in all dioecious freshwater mussel species lead to highly differentiated mitogenomes between the two sexes (Breton et al., 2010; Doucet-Beaupré et al., 2010). The female form has an extra protein coding gene called the F-ORF, while the male form shows some rearrangements of the gene order and features a unique 3' coding extension of the *cox2* subunit.

In the metazoan key phylum Placozoa (Schierwater, 2005) the mitochondrial genomes display several ancestral metazoan features (Dellaporta et al., 2006; Signorovitch et al., 2007). The extended genome size of up to 43 kb, the presence of introns, large intergenic regions and several ORFs of unknown function are shared characteristics with known metazoan outgroups like choanoflagellates such as *Monosiga brevicollis* (Burger et al., 2003) and therefore highlight a basal position of Placozoa at the root of the Metazoa. An unorthodox, fragmented *cox1* gene is assembled by trans-splicing involving split group I introns and requires U-to-C editing (Burger et al., 2009). While such features are not uncommon in other eukaryotic kingdoms, they are highly exceptional among Metazoa.

In sponges (Phylum Porifera), large-scale mitochondrial genome sequencing has given important insights into phylogenetic relationships in Demospongiae (Wang and Lavrov, 2008) and Homoscleromorpha (Gazave et al., 2010) and highlighted the value of mitochondrial data for phylogenetic inferences. With a size of up to 29 kb (Lavrov, 2010) the largest sponge mitochondrial genome is smaller than the smallest placozoan mitogenome, displaying the general metazoan tendency for mitogenome size reduction due to lower proportions of non-coding regions. On the other hand, they feature unique non-coding repetitive sequences capable of forming hairpin structures (Erpenbeck et al., 2009).

In cnidarians the mitochondrial genome architecture is another and unique character to understand the evolution of mitogenomes within this phylum. Only the basal Anthozoa harbour circular mitochondrial genomes with a size of up to 21 kb (Medina et al., 2006), while the other cnidarian classes display linear chromosomes. In Hydrozoa and Cubozoa they are in addition fragmented into several pieces (Voigt et al., 2008; Kayal et al., 2012). The largest number so far, eight chromosomes, has been found in the cubozoan *Alatina moseri*. Inverted repeat sequences form the telomeres

in these genomes, which may have originated from transposons or plasmids (Nosek and Tomáška, 2003). In combination with the tendency to reduce the number of encoded tRNA genes the linear mitogenome architecture underlines the derived status of medusozoan mitogenomes (Ender and Schierwater, 2003). Several cnidarian mitogenomes harbour group I introns in their *nad5* and *cox1* genes (Beagley et al., 1996; Fukami et al., 2007).

Homing endonucleases have been reported in the group I introns found in Cnidaria as well as Porifera (Rot et al., 2006; Goddard et al., 2006). They are thought to play a role in the mobility and origin of these introns, which apparently arose by horizontal transfer from fungi (see e.g. Szitenberg et al., 2010). A single case of group II introns was reported for the *cox1* gene in the mitogenomes of an annelid worm (Vallès et al., 2008).

The probably most derived mitogenomes within basal metazoan phyla are found in the Ctenophora. With a size of around 11 kb they are among the smallest animal mitogenomes. Like in cnidarians, ctenophoran mitogenomes have substantially reduced or even completely lost their mitochondrial encoded tRNA genes. The rRNAs as well as the protein coding sequences are highly derived in the Ctenophora, often leading to extremely long branches in phylogenetic analyses (Pett et al., 2011; Kohn et al., 2011).

Several instances of +1 frameshifts have been reported. For instance, a frameshift in the *nad3* gene is conserved in the sauropsid mitogenomes (Mindell et al., 1998; Russell and Beckenbach, 2008). Another well-described case is the *cob* gene in the ant genus *Polyrhachis* (Beckenbach et al., 2005) and in oysters (Milbury and Gaffney, 2005).

2.3. Genetic codes

Several codons do not have their standard meaning in some or all animal mitochondria. Interestingly, the variations concern only codons for which the first two positions do not determine the amino acid, and codons with a purine at the third position have been reassigned, see Table 2.

The reassignment of codon appears to be intimately linked to the chemical modifications of the tRNAs (Santos et al., 2004). Indeed, most (and maybe all) tRNAs that decode the known non-universal codons in animal mitochondria are chemically modified, usually in their anticodon wobble position. Evolutionary scenarios that link the emergence of code variations and changes in tRNA modifications are reviewed in (Watanabe and Yokobori, 2011), see also (Abascal et al., 2012).

2.4. NUMTs: nuclear copies of mitochondrial DNA

The exchange of DNA between the nucleus and mitochondria (as well as chloroplasts) is a common phenomenon throughout the eukaryotes. Repair of double-stranded breaks by non-homologous end-joining allows the inclusion of organellar DNA. Early in

evolution this has led to the transfer of a large fraction of the mitogenomic information to the nuclear genome. The products of many of these genes are now re-imported while other genes of mitochondrial origin have acquired novel functions elsewhere (Kleine et al., 2009). Recent nuclear inserts of mitochondrial sequences, which occur at surprisingly high rate probably limited by the rates of double strand breaks (Hazkani-Covo et al., 2010), however, give rise to non-coding and typically non-functional sequences. Nevertheless, a few NUMTs have been identified as the cause of genetic diseases (Chen et al., 2005). A recent example shows that they might also be a source of new spliceosomal introns (Curtis and Archibald, 2010). NUMTs can form a noticeable fraction of the nuclear genome: In the human genome at least 400 kb (Hazkani-Covo and Graur, 2007), in *Nasonia* ~43 kb and in *Apis* over 230 kb of the nuclear DNA consists of NUMTs. In contrast, some genomes such as that of *Drosophila melanogaster* are nearly devoid of mitochondrial DNA (Viljakainen et al., 2010).

From a phylogenetic point of view, NUMTs are primarily an annoyance. In PCR-based approaches, unrecognised divergent NUMTs can introduce noise and unrecognised biases. In the DNA barcoding approaches, furthermore, they cause a systematic overestimate of the number of species (Song et al., 2008). Box 1 in (Hazkani-Covo et al., 2010) discusses in some detail artefactual results associated with unrecognised NUMTs. Nevertheless current genome annotations do not routinely provide NUMT data tracks. On the other hand, NUMTs have also been suggested as an interesting tool in primate phylogeny (Hazkani-Covo, 2009).

3. Replication

3.1. Strand biases

The two strands of mammalian mitochondrial genome have been found to differ in their nucleotide composition in an asymmetric manner. While one strand is rich in G the other is G poor. Since this difference is physically measurable (Anderson et al., 1981) the G rich strand is also termed heavy strand (H-strand) and the other light strand (L-strand). In particular one strand of mammalian mitogenomes has been reported to be GT rich (Reyes et al., 1998). In contrast (both strands) of the mitogenomes of Arthropoda are AT rich (Crease, 1999). But still the asymmetry favouring GT on one strand and AC on the other can be clearly observed in the 3rd codon position of protein coding sequences of most metazoan mitogenomes, in particular for fourfold degenerate sites. For several species, e.g. *Branchiostoma* (Cephalochordata), *Florumetra* (Echinodermata), *Katharina* (Mollusca), and several Arthropod taxa (e.g. *Tigriopus*, *Argiope*, and *Euscorpius*), mitogenomes have opposite values (Hassanin et al., 2005; Wei et al., 2010).

The cause of the asymmetry is accounted to an asymmetric mutation process favouring transitions over transversions (Hassanin et al., 2005; Brown and Simpson, 1982). The likely source of this are hydrolytic deaminations of A and C (leading to A → G and C → T mutations) on the H-strand which is prone to mutations when it is single stranded which happens during replication as well as during transcription, but to a lesser extent. This is consistent with the correlations found between the nucleotide frequencies as well as with the substitution rates and the time spent in a single stranded conformation during replication as measured by the distances to origin of the light strand O_L (see Reyes et al., 1998; Krishnan et al., 2004). For some species a reversed bias in the nucleotide composition is observed, which might be caused by an inversion of the control region (Hassanin et al., 2005; Kilpert and Podsiadlowski, 2006; Wei et al., 2010).

Table 2
Variations of the universal Genetic Code (UC) in Animal Mitochondria.

Codon	UC	Mod	Examples
UGA	STOP	Trp	All Metazoa
AUA	Ile	Met	Bilateria except Planaria and Hemichordata
AAA	Lys	Asn	Echinodermata, some Platyhelminthes
AGA	Arg	Ser	Most Bilateria
		Gly	Tunicata
		STOP	Vertebrata
AGG	Arg	Ser	Most Bilateria
		Ser/Lys	Arthropoda
		Gly	Tunicata
		STOP	Vertebrata
UAA	STOP	Tyr	Planaria, Nematoda

Since nucleotide composition is related to amino acid composition (Min and Hickey, 2007) it has to be regarded in comparative analyses of amino acid sequences (Bernt et al., 2012a, in this special issue).

3.2. Model(s) of replication

The mtDNA is synthesised solely by DNA polymerase γ (Bolden et al., 1977). So far, three additional accessory factors are known to be involved in mtDNA synthesis. A single-stranded DNA-binding protein (mtSSB) has been reported (Tiranti et al., 1993), which is thought to bind to the single stranded parental H-strand to protect it. Furthermore mtDNA helicase activity has been reported for both vertebrates (bovine brain) (Hehman and Hauswirth, 1992) and invertebrates (sea urchin) (Roberti et al., 1996), and topoisomerases responsible for relaxing (type I) and introducing (type II) supercoils are involved in mtDNA replication (Kosovsky and Soslaw, 1993; Castora and Simpson, 1979).

The most widely accepted “orthodox” hypothesis explaining the mode of mtDNA replication is the so called *strand displacement* model (Robberson et al., 1972; Goddard and Wolstenholme, 1980; Clayton, 1982), see Fig. 2. We can give here only a brief outline and refer to the reviews by Shadel and Clayton (1997) and Taanman (1999) for much more detailed description of the replication process.

Primers for mtDNA H-strand replication are generated by the same process starting light strand transcription at the light strand promoter (LSP) (Chang et al., 1985) (see further Section 4). Whether light strand transcription is initiated or an RNA–DNA hybrid, called R-loop, is formed, is determined within the region of the heavy strand origin (O_H), i.e. the part of the D-Loop region that harbours three conserved sequence blocks (CSB) (Walberg and Clayton, 1981). The RNA in the R-loop is further processed by RNase MRP which yields the primers for mtDNA H-strand replication. The H-strand synthesis is then started by the elongation of the RNA primer by DNA Polymerase γ . Most DNA H-strand replication processes are terminated immediately after initiation at a location

harbouring one or more short (≈ 15 nt) termination associated sequences (TASs) and produce a stable triple stranded DNA–DNA hybrid called the D-loop structure. Alternatively, the H-strand replication proceeds unidirectionally from the short stretch of displaced mtDNA across the D-loop until it reaches the light-strand origin O_L . At this short (≈ 30 nt) non-coding region (which in Mammalia is located within a cluster of tRNAs), the displacement of the H-strand is thought to trigger the formation of a stem-loop structure (Martens and Clayton, 1979). This structure seems to be involved in the recognition by a DNA primase (Wong and Clayton, 1986) initiating L-strand replication. Since the stem loop structure is not universally present also the structural elements of the surrounding tRNAs might be involved in this process (Shadel and Clayton, 1997). L-strand replication then proceeds in the direction opposite of the direction of H-strand replication. The replication of both strands continues until the replication of the H-strand terminates and the two molecules are segregated. The remaining, so far unsynthesised part between O_H and O_L of the L-strand is finished after the segregation. Due to the delayed start of the L-strand replication the parental H-strand becomes single stranded with the advancing H-strand replication until the L-strand replication renders it double stranded again. The time which a position on the H-strand is single stranded depends on the distance to the origins as well as the speed of L- and H-strand replication. While in the mitogenomes of Mammalia the O_L is reported approximately 2/3 of the genome length after the O_H , this is even more extreme in insects, where it is located 97% after the O_H (see Saito et al., 2005). Replication of insect mtDNA is extremely asymmetric. Since the structure of the control region in mammals/chordates (Sbisá et al., 1997; Saccone et al., 1991) and insects (Saito et al., 2005; Zhang and Hewitt, 1997) is very different it is unclear at present whether the same mechanisms are responsible for initiation and termination of replication throughout the Metazoa.

As an alternative to the strand displacement model, a coupled mode of leading and lagging strand replication, called *strand-coupled* model, is discussed in the literature (Holt et al., 2000; Yang et al., 2002). Contrary to the orthodox model it is assumed that a




Fig. 2. Representation of the strand-displacement model of replication; mitochondrial genome with replication origins \approx as in human (A); H-strand replication has been initiated at O_H and proceeds towards O_L (A and B); L-strand replication has been started (D); H-strand replication finishes and segregation (E); the unfinished daughter replicate finished L-strand replication (F); light and heavy strand are distinguished by line width; H-Strand replication is indicated by a green arrow and L-strand replication by a blue arrow; single stranded regions of the H-strand are highlighted in red. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

replication bubble originates preferentially from an interval located near O_H spanning approximately one quarter of the mitogenome (Bowmaker et al., 2003; Reyes et al., 2005).

In this model stretches of RNA/ribonucleotides are unevenly incorporated mainly in the daughter L-strand (see also Pohjoismäki et al., 2011). It has been argued that different analysis methods may degrade these stretches resulting in replication intermediates with single stranded DNA that gave rise to the strand displacement model. Most recently, the possibility of bidirectional replication and multiple replication origins scattered throughout the mitogenome have been discussed for the strand-coupled model of replication (Bowmaker et al., 2003; Reyes et al., 2005). It is still unsolved whether there are two different replication mechanisms generating the different kinds of observed replication intermediates, if the replication mechanisms are employed in a tissue specific manner, or if one of the two models is wrong. For critical discussions of these issues see e.g. Clayton (2003); Fish et al. (2004); Brown et al. (2005); Pohjoismäki and Goffart (2011).

Not much is known about the replication of the mitogenomes with a non-circular chromosome organisation, see Section 2.2. In the case of *Globodera* it has been suggested that repetitive sequences present on each of the mini circles play a role in replication (Armstrong et al., 2000). In analogy to fungal mtDNA, individual circular chromosomes might replicate independently (Armstrong et al., 2000). The replication mechanism of the linear mitogenomes of Cnidarians remains unknown at present. Putative genomic regions involved in replication have been identified for *Hydra*, although several plausible mechanisms known for other genetic systems have been ruled out for this species (Voigt et al., 2008), see also (Kayal et al., 2012). It seems to be promising to evaluate alternative mechanisms for mitogenome replication known for non-metazoans with linear or multipartite mitogenomes (see e.g. Bendich, 2010, for some recent progress).

3.3. Mutation and repair

With a few exceptions such as Octocoralia (McFadden et al., 2006), metazoan mitogenomes are subject to strong mutational pressures (Brown et al., 1979) that are associated with an inaccurate DNA repair system compared to nuclear DNA (Bogenhagen, 1999; Gredilla, 2011), the lack of protective chromosome-associated proteins, and the replication mechanisms outlined above that expose single-stranded intermediates. In addition, mitochondrial DNA is subject to high levels of oxidative damage (Harman, 1972). In conjunction with relatively small population sizes and the virtual absence of recombination one would predict that mtDNA is subject to Muller's Ratchet (Muller, 1964), i.e. a gradual accumulation of deleterious mutations and eventually complete loss of function. The strength of this effect depends on population size. Indeed, an increased number of amino acid substitutions has been reported from large mammals, which tend to have small populations (Popadin et al., 2007). Overall, however, the function of mitochondrial proteins is preserved by very strong selection, reviewed in detail in (Castellana et al., 2011).

Positive selection has been reported in particular for vertebrates and to an even larger extent for insect mitogenomes (Bazin et al., 2006). These findings have cast doubt on the usefulness of mtDNA as a means for estimating effective population sizes (see e.g. Meiklejohn et al., 2007).

3.4. Genome rearrangements

The arrangement of genes on the mitogenome shows a very high degree of variation across the Metazoa. A comparison of closely related species with different gene orders suggests that there are several types of "elementary" rearrangement events, see Fig. 3:

inversions (e.g. Asakawa et al., 1995), transpositions (e.g. Macey et al., 1997), inverse transpositions (e.g. Boore et al., 1998) (i.e. a transposition in which the re-inserted fragment is inverted), and tandem duplications followed by the random loss of one of the copied genes (Boore, 2000) (TDRL). Variants of the latter (Mueller and Boore, 2005) as well as tandem duplications in which the subsequent loss is not randomly distributed among the copies (Lavrov et al., 2002) have been discussed. Additionally changes in the gene content, i.e. deletions (Lavrov and Brown, 2001) or duplications (Zhong et al., 2008) and changes in chromosome organisation (see Section 2.2) have been reported for metazoan mitogenomes. Rearrangements can reveal unexpected high-level clades such as the subdivision of placozoan lineages in two different groups that probably represent higher taxonomic units (Eitel and Schierwater, 2010).

It is not entirely clear at present whether the rearrangement types deduced from a comparative analysis of mitogenomes correspond to distinct specific molecular mechanisms. For instance transpositions and non-tandem duplication random loss events can be interpreted as special cases of the TDRL model. It is also unresolved whether inverse transpositions are truly elementary operations or the composite of two subsequent steps. Conversely, several different molecular mechanisms might be at work to produce the same observable rearrangement event.

A variety of possible mechanisms causing duplications have been discussed in the literature (Boore, 2000), several of which are related to the replication process, such as slipped strand mispairing during replication (Broughton and Dowling, 1997), imprecise termination (Mueller and Boore, 2005) (where too early termination results in a deletion and too late in a duplication event), and other enzymatic errors causing e.g. erroneous identification of the origin of light-strand replication (Macey et al., 1997). In a study of recent duplications in geckos, slipped-strand mispairing was favoured (Fujita et al., 2007). Another potential route to TDRLs is the head-to-head or head-to-tail dimerisation of linearised monomeric mitogenomes (Lavrov et al., 2002). The head-to-head configuration would result in an inverted duplicate. Note that any mechanism that can result in a whole genome duplication, e.g. imprecise termination, might also account for head-to-tail dimerisation.

Illegitimate recombination may also cause changes in gene order. It provides a plausible explanation in particular for non-tandem duplicated segments (Mueller and Boore, 2005). Certain features of the mitogenomic sequence, such as small direct repeats (Macey et al., 1997), have been proposed to evoke rearrangements. Duplications might occur due to the formation of stem loop structures which are involved in replication (e.g. at the O_L), or during the maturation of transcripts, see Section 4.

As an alternative to (tandem) duplication (non)-random loss, transpositions could also be caused by a separate mechanism akin to the propagation of transposable elements (Calos and Miller, 1980; Macey et al., 1997).

Irrespective of the duplication mechanism random loss can easily be explained by the accumulation of deleterious mutations in one of the copies of duplicated genes. Selection pressure on the size of the mitogenomic DNA will then rapidly remove the non-functional pseudogene. Remnants of such processes have been detected for instance as non-coding regions at positions where the deleted genes would be expected (Arndt and Smith, 1998) or as pseudogenes (Macey et al., 1997). Recently a large scale study on tRNAs provided numerous examples of pseudogenisation supporting duplication and inverse duplication based mechanisms (Jühling et al., 2012a). The presence of multiple functional copies after a duplication can facilitate tRNA recruitment, i.e. the change of the identity of a tRNA due to point mutations in the anticodon. Several recruitment events affecting the mitochondrial leucine tRNAs have




Fig. 3. Types of elementary rearrangement events exemplified for an artificial arrangement of five genes; from left to right: inversion, transposition, inverse transposition, tandem duplication random loss; pseudogenisation is represented by missing borders.

been reconstructed for the metazoan evolution (Higgs et al., 2003; Rawlings et al., 2003). Within the demosponges this phenomenon seems to be particularly frequent (Lavrov and Lang, 2005; Wang and Lavrov, 2011). This is phenomenon may lead to apparent rearrangements when tRNA identity and homology are used synonymously. This effect should be taken into account in the context of annotation (e.g. Dowton and Austin, 1999).

Inversions may be the result of intra-mitochondrial recombination (Dowton and Campbell, 2001; Lunt and Hyman, 1997). This mechanism may also lead to the formation of mini-circles (Dowton and Austin, 1999) and might also account for the heterogeneous gene contents observed in some clades, see Section 2.2. The mechanisms discussed above that can generate inverted duplicated segments, i.e. illegitimate recombination and head-to-head dimerisation, can be regarded as special cases producing apparent inversions of gene order.

Rearrangements often involve the replication origins (e.g. San Mauro et al., 2006), but also other hot spots of rearrangements have been described in the literature (Dowton and Austin, 1999). Based on the sparse species sampling available in early studies of mitogenomic rearrangements, tRNAs have been found to be involved more frequently in rearrangements than proteins and rRNAs (Wolstenholme, 1992; Macey et al., 1997). With the availability of a better species sampling and improved automatic methods for the comparison of mitochondrial gene orders (reviewed in Bernt et al., 2012a, in this special issue) large scale analyses on the properties of gene order evolution of mitogenomes have become possible. Case studies such as the manual analysis of Hymenoptera (Dowton et al., 2009) and two computational analyses of metazoan gene orders (Miklós and Hein, 2005; Bernt and Middendorf, 2011) confirm the conclusions of earlier studies and provide a more detailed quantitative picture of mitogenome rearrangements.

Less than 30% of the hymenopteran rearrangements are inversions, the majority being transpositions (Dowton et al., 2009). The 1:2 ratio of inversions to transpositions reported in (Miklós and Hein, 2005) may be due to the inclusion of the Deuterostomia or due to multiple counting of the same unique rearrangement. The extensive data set in (Bernt and Middendorf, 2011) allowed the most detailed analysis published so far. Slightly more than half of the reconstructed rearrangements have been found in non-chordates, but a considerable number of non-chordate rearrangements could not be reconstructed due to highly diverged gene orders. The detailed analysis of the reconstructed rearrangements in Protostomia in (Bernt and Middendorf, 2011) has shown 20% inversions, more than 55% transposition, and more than 10% each of inverse transpositions and TDRLs, respectively. Similar rearrangement frequencies have been observed in the reconstructed rearrangements of Metazoa. More than 3/4 of the rearrangements affect only tRNAs. In the vast majority of these cases, only a single tRNA is affected.

It is important to note that rearrangement rates are not only unevenly distributed within the genome, where clear hot spots can be identified, but also taxonomically non-uniform. While for some taxa, such as Chordata, only a few, similar gene orders are known, a different gene order is found for virtually every sequenced mitogenome in other clades. Multiple recent duplications have been described for some geckos (Fujita et al., 2007). The heavily rearranged gene orders of *Ascidia* form another counterexample to the generally preserved gene order of deuterostomes (Gissi et al., 2010).

Increased rearrangement rates were reported also for some insects (Shao et al., 2001) and within Mollusca (Grande et al., 2008). Genome rearrangement rates and nucleotide substitution frequency are correlated (Shao et al., 2003; Xu et al., 2006). It is unclear at present, which one is cause and which one is effect. Parasitic life styles seems to have an accelerating effect on the mutation rates (Castro et al., 2002).

4. Transcriptome

The mitochondrial genome is transcribed by a specialised machinery using the monomeric RNA polymerase POLRMT and specific mitochondrial transcription factors (mtTFA and the paralogous proteins mtTFB1 and mtTFB2). Interestingly, POLRMT also exhibits promoter-independent activity dependent on DNA supercoiling (Fukuoh et al., 2009). For reviews of the protein machinery involved in regulating mitochondrial gene expression we refer to Gagliardi et al. (2004); Shutt and Shadel (2010).

The primary transcripts are polycistronic precursors. Nuclear regulatory factors influence transcription levels both at the stage of initiation and termination (Asin-Cayuella and Gustafsson, 2007; Scarpulla, 2008). In mammals, the two alternative heavy strand promoters as well as the light strand promoter are located in the D-loop, reviewed e.g. by Asin-Cayuella and Gustafsson (2007) and Scarpulla (2008). They give rise to three primary transcripts. The HSP2 promoter within the *tRNA-Phe* generates a large transcript that terminates at the distal so-called D-TERM locus, containing a conserved A/T rich sequence motif immediately upstream of the *tRNA-Phe* gene (Camasamudram et al., 2003). The more abundant, shorter transcript starts at the HSP1 promoter upstream of the *tRNA-Phe* and encodes the two rRNAs and two tRNAs. It ends at proximal, so-called mt-TERM signal beyond the *rrnL* rRNA gene within the adjacent *tRNA-Leu* gene (Christianson and Clayton, 1986). Sequence box II, located in the D-loop, acts as a sequence-dependent termination element for transcription originating from the light strand promoter (Pham et al., 2006). This situation appears to be largely conserved throughout the vertebrates (Coucheron et al., 2011). The box II sequence itself is conserved at least throughout the mammals (Sbisá et al., 1997). In *Drosophila*, RNA synthesis starts at two promoters on the heavy (H) strand and three on the light (L) strand, largely matching the alternating blocks of genes located on heavy and light strand, respectively (Berthier et al., 1986; Torres et al., 2009). Large-scale rearrangements of gene order thus should be expected to affect also the structure of the mitochondrial transcriptome.

The primary transcripts are processed into mature rRNA and mRNA by cleavage of the 5' and 3' termini of the mitochondrial tRNAs. This processing, which is known as the RNA punctuation model (Ojala et al., 1980), is mediated by RNase P and tRNase Z endonucleases, respectively (Levinger et al., 2004). When multiple tRNA genes are present, as between *nad3* and *nad5* in *Drosophila*, they are removed sequentially from the 3' end (Stewart and Beckenbach, 2009). In both vertebrates and *Drosophila*, all mature mitochondrial mRNAs are monocistronic except the two bicistronic *nad4/nad4L* and *atp8/atp6* messages. Although *atp6* and *atp8* are found in adjacent positions in more than 90% of the sequenced mitogenomes, and in particular in nearly all Chordata

and Arthropoda, there are many exceptions, in particular in Mollusca, Cnidaria, Nematoda, and Platyhelminthes. The mRNAs are typically polyadenylated (Bobrowicz et al., 2008), although non-poly (A) mRNAs are also detectable (Mercer et al., 2011). An interesting feature of mitochondrial mRNAs is that many of the stop codons are complete only by polyadenylation (7 in human, 6 in cod, and at least 4 in fruit fly) so that they completely lack 3'UTRs (Nagaike et al., 2005; Stewart and Beckenbach, 2009). This appears to be a very frequent phenomenon without a distinctive phylogenetic distribution according to the RefSeq annotation.

cox1 features a 3'UTR complementary to the adjacent *tRNA-Ser*, and some mRNAs have heterogeneous 3'UTRs depending on the cell type (Temperley et al., 2010). This situation is conserved with minor differences at least among vertebrates (Coucheron et al., 2011). 5'UTRs are absent in general. Deep sequencing has been used already to generate detailed maps of transcript ends and processing sites (Lopez Sanchez et al., 2011; Brzezniak et al., 2011). Results demonstrated that mRNA abundance is heavily regulated by post-transcriptional processes (Torres et al., 2009; Mercer et al., 2011) and established the presence of non-canonical processing products. These appear to be linked in particular to the secondary structure of the tRNAs since non-canonical processing sites are associated with clusters of RNA secondary structures within the mRNAs (Mercer et al., 2011).

Mitochondrial rRNAs are also polyadenylated or at least feature a few non-encoded adenosines – as in the case of the *rrnS* of *Drosophila* (Stewart and Beckenbach, 2009). Like bacterial rRNAs, they are modified by both methylation and pseudouridylation at highly conserved positions (Ofengand and Bakin, 1997). All mitochondrial tRNAs require the addition of CCA to their 3' terminus by the mitochondrial tRNA-nucleotidyl transferase (Nagaike et al., 2001). Then they are chemically modified to ensure their proper folding, recognition, and base-pairing (Helm and Attardi, 2004; Messmer et al., 2009). Modifications of the anticodon stem, such as the hypermodified nucleoside 5-methylaminomethyl-2-thiouridylate at position 34, create the “wobble-base” necessary to enable the recognition of multiple codons (Agris et al., 2007).

In addition to the canonical genes a variety of non-coding RNA species are present in mitochondria (Lung et al., 2006). Two distinctive species of short RNAs are produced downstream of the tRNA 5' and 3' processing sites (Mercer et al., 2011). A small control region specific polyadenylated RNA species (7S RNA) with unknown function (Ojala et al., 1981) originates from the L-strand primary transcript. Rackham et al. (2011) report three long non-coding RNAs (lncRNAs) in mouse mitochondria that appear to have a role in the regulation of mitochondrial gene expression. Regulated antisense transcripts as well as a curious inverted repeat covalently linked to the *rrnL* RNA were reported to distinguish normal from tumour cells in human (Burzio et al., 2009). The mitochondrial transcription machinery, finally, also generates the RNA primer needed for initiation of heavy strand DNA synthesis. To this end, in mammals, a G-quadruplex structure is utilised to prematurely terminate transcription from the L-strand promoter (Wan-rooij et al., 2010).

5. Concluding remarks

Since the characterisation of the human mitochondrial genome in 1981 further genome sequencing led to the fall of several dogmas concerning the uniformity of animal mitochondrial genomes. At present the large set of observations is consistent with the idea that marked gene loss and mtDNA compaction occurred during the emergence of multi cellular animals, but the pronounced compaction was not coincident with the origin of the Metazoa. The best living surrogate for the ancestral metazoan mitochondrial genome

is the placozoan genomes, which combine both, characteristics from protozoans and metazoans. An evolutionary speciality is found in derived cnidarians, which have linearised their mitochondrial genomes. All bilaterian mitochondrial genomes are widely conserved with respect to size, gene structure and content. Particularly here the study of differences in gene arrangement and genetic codes provides a large, not yet fully explored repertoire of phylogenetic markers. Although mitochondrial genomes in diploblastic and triploblastic animals have taken different routes and likely evolved in parallel, their source as a rich repertoire of informative phylogenetic markers applies to both lineages.

Acknowledgments

This work was supported by the German Research Foundation (DFG) through the Project “Deep Metazoan Phylogeny” within SPP 1174 (Projects STA 850/3-2, STA 850/2).

References

- Abascal, F., Posada, D., Zardoya, R., 2012. The evolution of the mitochondrial genetic code in arthropods revisited. *Mitochondr. DNA* 23, 84–91.
- Agris, P.F., Vendeix, F.A., Graham, W.D., 2007. tRNA's wobble decoding of the genome: 40 years of modification. *J. Mol. Biol.* 366, 1–13.
- Alfonzo, J.D., Söll, D., 2009. Mitochondrial tRNA import—the challenge to understand has just begun. *Biol. Chem.* 390, 717–722.
- Anderson, S., Bankier, A.T., Barrell, B.G., de Bruijn, M.H.L., Coulson, A.R., Drouin, J., Eperon, I.C., Nierlich, D.P., Roe, B.A., Sanger, F., Schreier, P.H., Smith, A.J.H., Staden, R., Young, I.G., 1981. Sequence and organization of the human mitochondrial genome. *Nature* 290, 457–465.
- Armstrong, M.R., Blok, V.C., Phillips, M.S., 2000. A multipartite mitochondrial genome in the potato cyst nematode *Globodera pallida*. *Genetics* 154, 181–192.
- Arndt, A., Smith, M.J., 1998. Mitochondrial gene rearrangement in the sea cucumber genus *Cucumberia*. *Mol. Biol. Evol.* 15 (8), 9–16.
- Asakawa, S., Himeno, H., Miura, K., Watanabe, K., 1995. Nucleotide sequence and gene organization of the starfish *Asterina pectinifera* mitochondrial genome. *Genetics* 140 (3), 1047–1060.
- Asin-Cayuela, J., Gustafsson, C.M., 2007. Mitochondrial transcription and its regulation in mammalian cells. *Trends Biochem. Sci.* 32, 111–117.
- Bazin, E., Glémin, S., Galtier, N., 2006. Population size does not influence mitochondrial genetic diversity in animals. *Science* 312, 570–572.
- Beagley, C.T., Okada, N.A., Wolstenholme, D.R., 1996. Two mitochondrial group I introns in a metazoan, the sea anemone *Metridium senile*: one intron contains genes for subunits 1 and 3 of NADH dehydrogenase. *Proc. Natl. Acad. Sci. USA* 93 (11), 5619–5623.
- Beckenbach, A.T., Robson, S.K., Crozier, R.H., 2005. Single nucleotide +1 frameshifts in an apparently functional mitochondrial cytochrome b gene in ants of the genus *Polyrhachis*. *J. Mol. Evol.* 60, 141–152.
- Bendich, A.J., 2010. The end of the circle for yeast mitochondrial DNA. *Mol. Cell* 39, 831–832.
- Bernt, M., Braband, A., Middendorf, M., Misof, B., Rota-Stabelli, O., Stadler, P.F., 2012a. Bioinformatics methods for the comparative analysis of metazoan mitochondrial genome sequences. *Mol. Phylogenet. Evol.* <http://dx.doi.org/10.1016/j.ympev.2012.09.019>.
- Bernt, M., Donath, A., Jühling, F., Externbrink, F., Florentz, C., Fritzsche, G., Pütz, J., Middendorf, M., Stadler, P.F., 2012b. MITOS: improved de novo metazoan mitochondrial genome annotation. *Mol. Phylogenet. Evol.* <http://dx.doi.org/10.1016/j.ympev.2012.08.023>.
- Bernt, M., Middendorf, M., 2011. A method for computing an inventory of metazoan mitochondrial gene order rearrangements. *BMC Bioinform.* 12 (Suppl 9), S6.
- Berthier, F., Renaud, M., Alziari, S., Durand, R., 1986. RNA mapping on *Drosophila* mitochondrial DNA: precursors and template strands. *Nucl. Acids Res.* 14, 4519–4533.
- Bilewitch, J.P., Degnan, S.M., 2011. A unique horizontal gene transfer event has provided the octocoral mitochondrial genome with an active mismatch repair gene that has potential for an unusual self-contained function. *BMC Evol. Biol.* 11, 228.
- Bobrowicz, A.J., Lightowlers, R.N., Chrzanowska-Lightowlers, Z., 2008. Polyadenylation and degradation of mRNA in mammalian mitochondria: a missing link? *Biochem. Soc. Trans.* 36, 517–519.
- Bogenhagen, D.F., 1999. DNA repair 1999: repair of mtDNA in vertebrates. *Am. J. Hum. Genet.* 64, 1276–1281.
- Bogenhagen, D.F., Wang, Y., Shen, E.L., Kobayashi, R., 2003. Protein components of mitochondrial DNA nucleoids in higher eukaryotes. *Mol. Cell. Proteomics* 2, 1205–1216.
- Bolden, A., Noy, G.P., Weissbach, A., 1977. DNA polymerase of mitochondria is a gamma-polymerase. *J. Biol. Chem.* 252, 3351–3356.
- Boore, J.L., 1999. Animal mitochondrial genomes. *Nucl. Acids Res.* 27, 1767–1780.
- Boore, J.L., 2000. The duplication/random loss model for gene rearrangement exemplified by mitochondrial genomes of deuterostome animals. In: Sankoff,

- D., Nadeau, J.H. (Eds.), Comparative Genomics: Empirical and Analytical Approaches to Gene Order Dynamics, Map Alignment and the Evolution of Gene Families, Computational Biology Series, vol. 1. Kluwer Academic Publishers, pp. 133–147.
- Boore, J.L., 2006. The complete sequence of the mitochondrial genome of *Nautilus macromphalus* (Mollusca: Cephalopoda). *BMC Genom.* 7, 182.
- Boore, J.L., Lavrov, D.V., Brown, W.M., 1998. Gene translocation links insects and crustaceans. *Nature* 392, 667–668.
- Bourlat, S.J., Nielsen, C., Economou, A.D., Telford, M.J., 2008. Testing the new animal phylogeny: a phylum level molecular analysis of the animal kingdom. *Mol. Phylogenet. Evol.* 49, 23–31.
- Bowmaker, M., Yang, M.Y., Yasukawa, T., Reyes, A., Jacobs, H.T., Huberman, J.A., Holt, J.J., 2003. Mammalian mitochondrial DNA replicates bidirectionally from an initiation zone. *J. Biol. Chem.* 278 (51), 50961–50969.
- Braband, A., Cameron, S.L., Podsiadlowski, L., Daniels, S.R., Mayer, G., 2010. The mitochondrial genome of the onychophoran *Opisthopatus cinctipes* (Peripatopsidae) reflects the ancestral mitochondrial gene arrangement of Panarthropoda and Ecdysozoa. *Mol. Phylogenet. Evol.* 57, 285–292.
- Breton, S., Beaupré, H.D., Stewart, D.T., Piontkivska, H., Karmakar, M., Bogan, A.E., Blier, P.U., Hoeh, W.R., 2009. Comparative mitochondrial genomics of freshwater mussels (Bivalvia: Unionoida) with doubly uniparental inheritance of mtDNA: gender-specific open reading frames and putative origins of replication. *Genetics* 183 (4), 1575–1589.
- Breton, S., Stewart, D.T., Shephardson, S., Trdan, R.J., Bogan, A.E., Chapman, E.G., Ruminas, A.J., Piontkivska, H., Hoeh, W.R., 2010. Novel protein genes in animal mtDNA: a new sex determination system in freshwater mussels (Bivalvia: Unionoida)? *Mol. Biol. Evol.* 28, 1645–1659.
- Broughton, R.E., Dowling, T.E., 1997. Evolutionary dynamics of tandem repeats in the mitochondrial DNA control region of the minnow *Cyprinella spiloptera*. *Mol. Biol. Evol.* 14, 1187–1196.
- Brown, G.G., Simpson, M.V., 1982. Novel features of animal mtDNA evolution as shown by sequences of two rat cytochrome oxidase subunit II genes. *Proc. Natl. Acad. Sci. USA* 79, 3246–3250.
- Brown, T.A., Cecconi, C., Tkachuk, A.N., Bustamante, C., Clayton, D.A., 2005. Replication of mitochondrial DNA occurs by strand displacement with alternative light-strand origins, not via a strand-coupled mechanism. *Gene Dev.* 19, 2466–2476.
- Brown, W.M., George Jr., M., Wilson, A.C., 1979. Rapid evolution of animal mitochondrial DNA. *Proc. Natl. Acad. Sci. USA* 76, 1967–1971.
- Brzezniański, L.K., Bijata, M., Szczesny, R.J., Stepien, P.P., 2011. Involvement of human ELAC2 gene product in 3' end processing of mitochondrial tRNAs. *RNA Biol.* 8, 616–626.
- Burger, G., Forget, L., Zhu, Y., Gray, M.W., Lang, B.F., 2003. Unique mitochondrial genome architecture in unicellular relatives of animals. *Proc. Natl. Acad. Sci. USA* 100, 892–897.
- Burger, G., Yan, Y., Javadi, P., Lang, B.F., 2009. Group I-intron trans-splicing and mRNA editing in the mitochondria of placozoan animals. *Trends Genet.* 25, 381–386.
- Burzio, V.A., Villota, C., Villegas, J., Landerer, E., Boccardo, E., Villa, L.L., Martínez, R., Lopez, C., Gaete, F., Toro, V., Rodriguez, X., Burzio, L.O., 2009. Expression of a family of noncoding mitochondrial RNAs distinguishes normal from cancer cells. *Proc. Natl. Acad. Sci. USA* 106, 9430–9434.
- Calos, M.P., Miller, J.H., 1980. Transposable elements. *Cell* 20, 579–595.
- Camasmudram, V., Fang, J.K., Avadhani, N.G., 2003. Transcription termination at the mouse mitochondrial H-strand promoter distal site requires an A/T rich sequence motif and sequence specific DNA binding proteins. *Eur. J. Biochem.* 270, 1128–1140.
- Cameron, S.L., Yoshizawa, K., Mizukoshi, A., Whiting, M.F., Johnson, K.P., 2011. Mitochondrial genome deletions and minicircles are common in lice (insecta: Phthiraptera). *BMC Genom.* 12, 394.
- Castellana, S., Vicario, S., Saccone, C., 2011. Evolutionary patterns of the mitochondrial genome in Metazoa: exploring the role of mutation and selection in mitochondrial protein coding genes. *Genom. Biol. Evol.* 3, 1067–1079.
- Castora, F.J., Simpson, M.V., 1979. Search for a DNA gyrase in mammalian mitochondria. *J. Biol. Chem.* 254, 11193–11195.
- Castro, L.R., Austin, A.D., Dowton, M., 2002. Contrasting rates of mitochondrial molecular evolution in parasitic Diptera and Hymenoptera. *Mol. Biol. Evol.* 19 (7), 1100–1113.
- Chacinska, A., Koehler, C.M., Milenkovic, D., Lithgow, T., Pfanner, N., 2009. Importing mitochondrial proteins: machineries and mechanisms. *Cell* 138 (4), 628–644.
- Chang, D.D., Hauswirth, W.W., Clayton, D.A., 1985. Replication priming and transcription initiate from precisely the same site in mouse mitochondrial DNA. *EMBO J.* 4, 1559–1567.
- Chen, J.M., Chuzhanova, N., Stenson, P.D., Ferec, C., Cooper, D.N., 2005. Meta-analysis of gross insertions causing human genetic disease: novel mutational mechanisms and the role of replication slippage. *Hum. Mutat.* 25, 207–221.
- Christianson, T.W., Clayton, D.A., 1986. In vitro transcription of human mitochondrial DNA: accurate termination requires a region of DNA sequence that can function bidirectionally. *Proc. Natl. Acad. Sci. USA* 83, 6277–6281.
- Clayton, D., 2003. Mitochondrial DNA replication: what we know. *IUBMB Life* 55, 213–217.
- Clayton, D.A., 1982. Replication of animal mitochondrial DNA. *Cell* 28, 693–705.
- Coucheron, D.H., Nymark, M., Breines, R., Karlsen, B.O., Andreassen, M., Jørgensen, T.E., Moum, T., Johansen, S.D., 2011. Characterization of mitochondrial mRNAs in codfish reveals unique features compared to mammals. *Curr. Genet.* 57, 213–222.
- Crease, T.J., 1999. The complete sequence of the mitochondrial genome of *Daphnia pulex* (Cladocera: Crustacea). *Gene* 233 (1–2), 89–99.
- Curtis, B.A., Archibald, J.M., 2010. A spliceosomal intron of mitochondrial DNA origin. *Curr. Biol.* 20, R919–R920.
- Dellaporta, S.L., Xu, A., Sagasser, S., Jakob, W., Moreno, M.A., Buss, L.W., Schierwater, B., 2006. Mitochondrial genome of *Trichoplax adhaerens* supports Placozoa as the basal lower metazoan phylum. *Proc. Natl. Acad. Sci. USA* 103, 8751–8756.
- Dörner, M., Altmann, M., Pääbo, S., Mörl, M., 2001. Evidence for import of a lysyl-tRNA into marsupial mitochondria. *Mol. Biol. Cell* 12, 2688–2698.
- Doucet-Beaupré, H., Breton, S., Chapman, E.G., Blier, P.U., Bogan, A.E., Stewart, D.T., Hoeh, W.R., 2010. Mitochondrial phylogenomics of the Bivalvia (Mollusca): searching for the origin and mitogenomic correlates of doubly uniparental inheritance of mtDNA. *BMC Evol. Biol.* 10, 50.
- Dowton, M., Austin, A.D., 1999. Evolutionary dynamics of a mitochondrial rearrangement “hot spot” in the Hymenoptera. *Mol. Biol. Evol.* 16, 298–309.
- Dowton, M., Cameron, S.L., Dowavic, J.L., Austin, A.D., Whiting, M., 2009. Characterization of 67 mitochondrial tRNA gene rearrangements in the hymenoptera suggests that mitochondrial tRNA gene position is selectively neutral. *Mol. Biol. Evol.* 26, 1607–1617.
- Dowton, M., Campbell, N.J., 2001. Intramitochondrial recombination – is it why some mitochondrial genes sleep around? *Trends Ecol. Evol.* 16, 269–271.
- Eitel, M., Schierwater, B., 2010. The phylogeography of the Placozoa suggests a taxon rich phylum in tropical and subtropical waters. *Mol. Ecol.* 19, 2315–2327.
- Ender, A., Schierwater, B., 2003. Placozoa are not derived cnidarians: evidence from molecular morphology. *Mol. Biol. Evol.* 20, 130–134.
- Erpenbeck, D., Voigt, O., Wörheide, G., Lavrov, D., 2009. The mitochondrial genomes of sponges provide evidence for multiple invasions by Repetitive Hairpin-forming Elements (RHE). *BMC Genom.* 10, 591.
- Fernández-Silva, P., Enriquez, J.A., Montoya, J., 2003. Replication and transcription of mammalian mitochondrial DNA. *Exp. Physiol.* 88, 41–56.
- Fish, J., Raule, N., Attardi, G., 2004. Discovery of a major D-Loop replication origin reveals two modes of human mtDNA synthesis. *Science* 306, 2098–2101.
- Fujita, M.K., Boore, J.L., Moritz, C., 2007. Multiple origins and rapid evolution of duplicated mitochondrial genes in parthenogenetic geckos (*Heteronotia binoei*; Squamata, Gekkonidae). *Mol. Biol. Evol.* 24, 2775–2786.
- Fukami, H., Chen, C., Chiou, C.-Y., Knowlton, N., 2007. Novel group I introns encoding a putative homing endonuclease in the mitochondrial *cox1* gene of scleractinian corals. *J. Mol. Evol.* 64, 591–600.
- Fukuoh, A., Ohgaki, K., Hatae, H., Kuraoka, I., Aoki, Y., Uchiyama, T., Jacobs, H.T., Kang, D., 2009. DNA conformation-dependent activities of human mitochondrial RNA polymerase. *Genes Cells* 14, 1029–1042.
- Gagliardi, D., Stepien, P.P., Temperley, R.J., Lightowlers, R.N., Chrzanoska-Lightowlers, Z.M., 2004. Messenger RNA stability in mitochondria: different means to an end. *Trends Genet.* 20, 260–267.
- Gazave, E., Lapebie, P., Renard, E., Vacelet, J., Rocher, C., Ereskovsky, A.V., Lavrov, D.V., Borchellini, C., 2010. Molecular phylogeny restores the supra-generic subdivision of homoscleromorph sponges (Porifera, Homoscleromorpha). *PLoS ONE* 5, e14290.
- Gissi, C., Iannelli, F., Pesole, G., 2008. Evolution of the mitochondrial genome of metazoa as exemplified by comparison of congeneric species. *Heredity* 101, 301–320.
- Gissi, C., Pesole, G., Mastrototaro, F., Iannelli, F., Guida, V., Griggio, F., 2010. Hypervariability of ascidian mitochondrial gene order: exposing the myth of deuterostome organelle genome stability. *Mol. Biol. Evol.* 27 (2), 211–215.
- Goddard, J.M., Wolstenholme, D.R., 1980. Origin and direction of replication in mitochondrial DNA molecules from the genus *Drosophila*. *Nucl. Acids Res.* 8, 741–757.
- Goddard, M.R., Leigh, J., Roger, A.J., Pemberton, A.J., 2006. Invasion and persistence of a selfish gene in the Cnidaria. *PLoS ONE* 1, e3.
- Grande, C., Templado, J., Zardoya, R., 2008. Evolution of gastropod mitochondrial genome arrangements. *BMC Evol. Biol.* 8, 61.
- Gredilla, R., 2011. DNA damage and base excision repair in mitochondria and their role in aging. *J. Aging Res.* 2011, 257093.
- Haen, K.M., Pett, W., Lavrov, D.V., 2010. Parallel loss of nuclear-encoded mitochondrial aminoacyl-tRNA synthetases and mtDNA-encoded tRNAs in Cnidaria. *Mol. Biol. Evol.* 27, 2216–2219.
- Harman, D., 1972. A biologic clock: the mitochondria? *J. Am. Geriatr. Soc.* 20, 145–147.
- Hassanin, A., Lger, N., Deutsch, J., 2005. Evidence for multiple reversals of asymmetric mutational constraints during the evolution of the mitochondrial genome of Metazoa, and consequences for phylogenetic inferences. *Syst. Biol.* 54, 277–298.
- Hazkani-Covo, E., 2009. Mitochondrial insertions into primate nuclear genomes suggest the use of numts as a tool for phylogeny. *Mol. Biol. Evol.* 26, 2175–2179.
- Hazkani-Covo, E., Graur, D., 2007. A comparative analysis of numt evolution in human and chimpanzee. *Mol. Biol. Evol.* 24, 13–18.
- Hazkani-Covo, E., Zeller, R., Martin, W., 2010. Molecular poltergeists: Mitochondrial DNA copies (numts) in sequenced nuclear genomes. *PLoS Genet.* 6, e1000834.
- Helman, G.L., Hauswirth, W.W., 1992. DNA helicase from mammalian mitochondria. *Proc. Natl. Acad. Sci. USA* 89, 8562–8566.
- Helfenbin, K.G., Fourcade, H.M., Vanjani, R.G., Boore, J.L., 2004. The mitochondrial genome of *Paraspadella gotoi* is highly reduced and reveals that chaetognaths are a sister group to protostomes. *Proc. Natl. Acad. Sci. USA* 101, 10639–10643.

- Helm, M., 2006. Post-transcriptional nucleotide modification and alternative folding of RNA. *Nucl. Acids Res.* 34, 721–733.
- Helm, M., Attardi, G., 2004. Nuclear control of cloverleaf structure of human mitochondrial tRNA^{Leu}. *J. Mol. Biol.* 337, 545–560.
- Higgs, P.G., Jameson, D., Jow, H., Ratray, M., 2003. The evolution of tRNA-Leu genes in animal mitochondrial genomes. *J. Mol. Evol.* 57, 435–445.
- Holt, I.J., Lorimer, H.E., Jacobs, H.T., 2000. Coupled leading- and lagging-strand synthesis of mammalian mitochondrial DNA. *Cell* 100, 515–524.
- Huynen, M., 2010. Mosaic origin of the mitochondrial proteome. *Proteomics* 10, 4012–4024.
- Jühling, F., Pütz, J., Bernt, M., Donath, A., Middendorf, M., Florentz, C., Stadler, P.F., 2012a. Improved systematic tRNA gene annotation allows new insights into the evolution of mitochondrial tRNA structures and into the mechanisms of mitochondrial genome rearrangements. *Nucl. Acids Res.* 40, 2833–2845.
- Jühling, F., Pütz, J., Florentz, C., Stadler, P.F., 2012b. Armless mitochondrial tRNAs in *Neopleura* (Nematoda). *RNA Biol.* 9, 1161–1166.
- Kayal, E., Bentlage, B., Collins, A.G., Kayal, M., Pirro, S., Lavrov, D.V., 2012. Evolution of linear mitochondrial genomes in medusozoan cnidarians. *Genome Biol. Evol.* 4, 1–12.
- Kayal, E., Lavrov, D.V., 2008. The mitochondrial genome of *Hydra oligactis* (Cnidaria, Hydrozoa) sheds new light on animal mtDNA evolution and cnidarian phylogeny. *Gene* 410, 177–186.
- Kilpert, F., Podsiadlowski, L., 2006. The complete mitochondrial genome of the common sea slater, *Ligia oceanica* (Crustacea, Isopoda) bears a novel gene order and unusual control region features. *BMC Genom.* 7, 241.
- Kleine, T., Maier, U.G., Leister, D., 2009. DNA transfer from organelles to the nucleus: the idiosyncratic genetics of endosymbiosis. *Annu. Rev. Plant Biol.* 60, 115–138.
- Kohn, A.B., Citarella, M.R., Kocot, K.M., Bobkova, Y.V., Halanych, K.M., Moroz, L.L., 2011. Rapid evolution of the compact and unusual mitochondrial genome in the ctenophore, *Pleurobrachia bachei*. *Mol. Phylog. Evol.* 63, 203–207.
- Kosovsky, M.J., Soslau, G., 1993. Immunological identification of human platelet mitochondrial DNA topoisomerase I. *Biochim. Biophys. Acta* 1164, 101–107.
- Krishnan, N.M., Seligmann, H., Raina, S.Z., Pollock, D.D., 2004. Detecting gradients of asymmetry in site-specific substitutions in mitochondrial genomes. *DNA Cell Biol.* 23, 707–714.
- Lavrov, D.V., 2010. Rapid proliferation of repetitive palindromic elements in mtDNA of the endemic baikalian sponge *Lubomirskia baicalensis*. *Mol. Biol. Evol.* 27, 757–760.
- Lavrov, D.V., Boore, J.L., Brown, W.M., 2002. Complete mtDNA sequences of two millipedes suggest a new model for mitochondrial gene rearrangements: duplication and nonrandom loss. *Mol. Biol. Evol.* 19, 163–169.
- Lavrov, D.V., Brown, W.M., 2001. Trichinella spiralis mtDNA: a nematode mitochondrial genome that encodes a putative *ATP8* and normally structured tRNAs and has a gene arrangement relatable to those of coelomate metazoans. *Genetics* 157, 621–637.
- Lavrov, D.V., Lang, B.F., 2005. Transfer RNA gene recruitment in mitochondrial DNA. *Trends Genet.* 21, 129–133.
- Levinger, L., Mörl, M., Florentz, C., 2004. Mitochondrial tRNA 3' end metabolism and human disease. *Nucl. Acids Res.* 32, 5430–5441.
- Lopez Sanchez, M.L., Mercer, T.R., Davies, S.M., Shearwood, A.M., Nygård, K.K., Richman, T.R., Mattick, J.S., Rackham, O., Filipovska, A., 2011. RNA processing in human mitochondria. *Cell Cycle* 10, 2904–2916.
- Lung, B., Zemann, A., Madej, M.J., Schuelke, M., Techritz, S., Ruf, S., Bock, R., Hüttenhofer, A., 2006. Identification of small non-coding RNAs from mitochondria and chloroplasts. *Nucl. Acids Res.* 34, 3842–3852.
- Lunt, D.H., Hyman, B.C., 1997. Animal mitochondrial DNA recombination. *Nature* 387, 247.
- Macey, J.R., Larson, A., Ananjeva, N.B., Fang, Z., Papenfuss, T.J., 1997. Two novel gene orders and the role of light-strand replication in rearrangement of the vertebrate mitochondrial genome. *Mol. Biol. Evol.* 14, 91–104.
- Martens, P.A., Clayton, D.A., 1979. Mechanism of mitochondrial DNA replication in mouse L-cells: localization and sequence of the light-strand origin of replication. *J. Mol. Biol.* 135 (2), 327–351.
- McBride, H.M., Neuspiel, M., Wasiak, S., 2006. Mitochondria: more than just a powerhouse. *Curr. Biol.* 16, R551–R560.
- McFadden, C.S., France, S.C., Sánchez, J.A., Alderslade, P., 2006. A molecular phylogenetic analysis of the *Octocorallia* (Cnidaria:Anthozoa) based on mitochondrial protein-coding sequences. *Mol. Phylog. Evol.* 41, 513–527.
- Medina, M., Collins, A.G., Takaoka, T.L., Kuehl, J.V., Boore, J.L., 2006. Naked corals: Skeleton loss in *Scleractinia*. *Proc. Natl. Acad. Sci. USA* 103, 9096–9100.
- Meiklejohn, C.D., Montooth, K.L., Rand, D.M., 2007. Positive and negative selection on the mitochondrial genome. *Trends Genet.* 23, 259–263.
- Mercer, T.R., Neph, S., Dinger, M.E., Crawford, J., Smith, M.A., Shearwood, A.M., Haugen, E., Bracken, C.P., Rackham, O., Stamatoyannopoulos, J.A., Filipovska, A., Mattick, J.S., 2011. The human mitochondrial transcriptome. *Cell* 146, 645–658.
- Messmer, M., Pütz, J., Suzuki, T., Sauter, C., Sissler, M., Florentz, C., 2009. Tertiary network in mammalian mitochondrial tRNA^{Asp} revealed by solution probing and phylogeny. *Nucl. Acids Res.* 37, 6881–6895.
- Miklós, I., Hein, J., 2005. Genome rearrangement in mitochondria and its computational biology. In: *Comparative Genomics, RECOMB 2005 International Workshop, RCG 2005, Proceedings. Lecture Notes in Computer Science*, vol. 3388. Springer, pp. 85–96.
- Milbury, C., Gaffney, P., 2005. Complete mitochondrial DNA sequence of the eastern oyster *Crassostrea virginica*. *Mar. Biotechnol.* 7, 697–712.
- Milbury, C.A., Lee, J., Cannone, J.J., Gaffney, P.M., Gutell, R.R., 2010. Fragmentation of the large subunit ribosomal RNA gene in oyster mitochondrial genomes. *BMC Genom.* 11, 485.
- Min, X.J., Hickey, D.A., 2007. DNA asymmetric strand bias affects the amino acid composition of mitochondrial proteins. *DNA Res.* 14 (5), 201–206.
- Mindell, D.P., Sorenson, M.D., Dimcheff, D.E., 1998. An extra nucleotide is not translated in mitochondrial *ND3* of some birds and turtles. *Mol. Biol. Evol.* 15, 1568–1571.
- Mueller, R.L., Boore, J.L., 2005. Molecular mechanisms of extensive mitochondrial gene rearrangement in plethodontid salamanders. *Mol. Biol. Evol.* 22, 2104–2112.
- Muller, H.J., 1964. The relation of recombination to mutational advance. *Mutat. Res.* 1, 2–9.
- Nagaike, T., Suzuki, T., Katoh, T., Ueda, T., 2005. Human mitochondrial mRNAs are stabilized with polyadenylation regulated by mitochondria-specific poly(A) polymerase and polynucleotide phosphorylase. *J. Biol. Chem.* 280, 19721–19727.
- Nagaike, T., Suzuki, T., Tomari, Y., Takemoto-Hori, C., Negayama, F., Watanabe, K., Ueda, T., 2001. Identification and characterization of mammalian mitochondrial tRNA nucleotidyltransferases. *J. Biol. Chem.* 276, 40041–40049.
- Nosek, J., Tomáška, L., 2003. Mitochondrial genome diversity: evolution of the molecular architecture and replication strategy. *Curr. Genet.* 44, 73–84.
- Ofengand, J., Bakin, A., 1997. Mapping to nucleotide resolution of pseudouridine residues in large subunit ribosomal RNAs from representative eukaryotes, prokaryotes, archaeobacteria, mitochondria and chloroplasts. *J. Mol. Biol.* 266, 246–268.
- Ojala, D., Crews, S., Montoya, J., Gelfand, R., Attardi, G., 1981. A small polyadenylated RNA (7 S RNA), containing a putative ribosome attachment site, maps near the origin of human mitochondrial DNA replication. *J. Mol. Biol.* 150, 303–314.
- Ojala, D., Merkel, C., Gelfand, R., Attardi, G., 1980. The tRNA genes punctuate the reading of genetic information in human mitochondrial DNA. *Cell* 22, 393–403.
- Pett, W., Ryan, J.F., Pang, K., Mullikin, J.C., Martindale, M., Baxeveanis, A.D., Lavrov, D.V., 2011. Extreme mitochondrial evolution in the ctenophore *Mnemiopsis leidyi*: insight from mtDNA and the nuclear genome. *Mitochondrial DNA* 22, 130–142.
- Pham, X.H., Farge, G., Shi, Y., Gaspari, M., Gustafsson, C.M., Falkenberg, M., 2006. Conserved sequence box II directs transcription termination and primer formation in mitochondria. *J. Biol. Chem.* 281, 24647–24652.
- Podsiadlowski, L., Brabant, A., Mayer, G., 2008. The complete mitochondrial genome of the onychophoran *Epiperipatus bioleyi* reveals a unique transfer RNA set and provides further support for the ecdysozoa hypothesis. *Mol. Biol. Evol.* 25, 42–51.
- Pohjoismäki, J.L.O., Goffart, S., 2011. Of circles, forks and humanity: topological organisation and replication of mammalian mitochondrial DNA. *BioEssays* 33, 290–299.
- Pohjoismäki, J.L.O., Holmes, J.B., Wood, S.R., Yang, M.-Y., Yasukawa, T., Reyes, A., Bailey, L.J., Cluett, T.J., Goffart, S., Willcox, S., Rigby, R.E., Jackson, A.P., Spellbrink, J.N., Griffith, J.D., Crouch, R.J., Jacobs, H.T., Holt, I.J., 2011. Mammalian mitochondrial DNA replication intermediates are essentially duplex, but contain extensive tracts of RNA/DNA hybrid. *J. Mol. Biol.* 397, 1144–1155.
- Popadin, K., Polishchuk, L.V., Mamirova, L., Knorre, D., Gunbin, K., 2007. Accumulation of slightly deleterious mutations in mitochondrial protein-coding genes of large versus small mammals. *Proc. Natl. Acad. Sci. USA* 104, 13390–13395.
- Rackham, O., Shearwood, A.M., Mercer, T.R., Davies, S.M., Mattick, J.S., Filipovska, A., 2011. Long noncoding RNAs are generated from the mitochondrial genome and regulated by nuclear-encoded proteins. *RNA* 17, 2085–2093.
- Raimond, R., Marcadé, I., Bouchon, D., Rigaud, T., Bossy, J.-P., Souty-Grosset, C., 1999. Organization of the large mitochondrial genome in the isopod *Armadiidium vulgare*. *Genetics* 151, 203–210.
- Rawlings, T.A., Collins, T.M., Bieler, R., 2003. Changing identities: tRNA duplication and remodeling within animal mitochondrial genomes. *Proc. Natl. Acad. Sci. USA* 100, 15700–15705.
- Reyes, A., Gissi, C., Pesole, G., Saccone, C., 1998. Asymmetrical directional mutation pressure in the mitochondrial genome of mammals. *Mol. Biol. Evol.* 15, 957–966.
- Reyes, A., Yang, M.Y., Bowmaker, M., Holt, I.J., 2005. Bidirectional replication initiates at sites throughout the mitochondrial genome of birds. *J. Biol. Chem.* 280, 3242–3250.
- Robberson, D.L., Kasamatsu, H., Vinograd, J., 1972. Replication of mitochondrial DNA. Circular replicative intermediates in mouse L cells. *Proc. Natl. Acad. Sci. USA* 69 (3), 737–741.
- Roberti, M., Musicco, C., Polosa, P.L., Gadaleta, M.N., Cantatore, P., 1996. DNA-helicase activity from sea urchin mitochondria. *Biochem. Biophys. Res. Commun.* 219, 134–139.
- Rot, C., Goldfarb, I., Ilan, M., Huchon, D., 2006. Putative cross-kingdom horizontal gene transfer in sponge (Porifera) mitochondria. *BMC Evol. Biol.* 6, 71.
- Russell, R., Beckenbach, A., 2008. Recoding of translation in turtle mitochondrial genomes: programmed frameshift mutations and evidence of a modified genetic code. *J. Mol. Evol.* 67, 682–695.
- Saccone, C., Pesole, G., Sbisá, E., 1991. The main regulatory region of mammalian mitochondrial DNA: structure-function model and evolutionary pattern. *J. Mol. Evol.* 33, 83–91.
- Saito, S., Tamura, K., Aotsuka, T., 2005. Replication origin of mitochondrial DNA in insects. *Genetics* 171, 1695–1705.

- San Mauro, D., Gower, D.J., Zardoya, R., Wilkinson, M., 2006. A hotspot of gene order rearrangement by tandem duplication and random loss in the vertebrate mitochondrial genome. *Mol. Biol. Evol.* 23, 227–234.
- Santos, M.A.S., Moura, G., Massey, S.E., Tuite, M.F., 2004. Driving change: the evolution of alternative genetic codes. *Trends Genet.* 20 (2), 95–102.
- Satta, Y., Ishiwa, H., Chigusa, S.I., 1987. Analysis of nucleotide substitutions of mitochondrial DNAs in *Drosophila melanogaster* and its sibling species. *Mol. Biol. Evol.* 4, 638–650.
- Sbisá, E., Tanzariello, F., Reyes, A., Pesole, G., Saccone, C., 1997. Mammalian mitochondrial D-loop region structural analysis: identification of new conserved sequences and their functional and evolutionary implications. *Gene* 205, 125–140.
- Scarpulla, R.C., 2008. Transcriptional paradigms in mammalian mitochondrial biogenesis and function. *Physiol. Rev.* 88, 611–638.
- Schierwater, B., 2005. My favorite animal, *Trichoplax adhaerens*. *BioEssays* 27, 1294–1302.
- Segovia, R., Pett, W., Treweek, S., Lavrov, D.V., 2011. Extensive and evolutionarily persistent mitochondrial tRNA editing in velvet worms (phylum Onychophora). *Mol. Biol. Evol.* 28, 2873–2881.
- Shadel, G.S., Clayton, D.A., 1997. Mitochondrial DNA maintenance in vertebrates. *Annu. Rev. Biochem.* 66, 409–435.
- Shao, R., Barker, S.C., 2011. Chimeric mitochondrial minichromosomes of the human body louse, *Pediculus humanus*: evidence for homologous and non-homologous recombination. *Gene* 473, 36–43.
- Shao, R., Campbell, N.J.H., Schmidt, E.R., Barker, S.C., 2001. Increased rate of gene rearrangement in the mitochondrial genomes of three orders of hemipteroid insects. *Mol. Biol. Evol.* 18 (9), 1828–1832.
- Shao, R., Dowton, M., Murrell, A., Barker, S.C., 2003. Rates of gene rearrangement and nucleotide substitution are correlated in the mitochondrial genomes of insects. *Mol. Biol. Evol.* 20 (10), 1612–1619.
- Shao, R., Kirkness, E.F., Barker, S.C., 2009. The single mitochondrial chromosome typical of animals has evolved into 18 minichromosomes in the human body louse, *Pediculus humanus*. *Genome Res.* 19, 904–912.
- Shutt, T.E., Shadel, G.S., 2010. Inventory of the human mitochondrial gene expression machinery with links to disease. *Environ. Mol. Mutagen.* 51, 360–379.
- Signorovitch, A.Y., Buss, L.W., Dellaporta, S.L., 2007. Comparative genomics of large mitochondria in placozoans. *PLoS Genet.* 3, e13.
- Song, H., Buhay, J.E., Whiting, M.F., Crandall, K.A., 2008. Many species in one: DNA barcoding overestimates the number of species when nuclear mitochondrial pseudogenes are coamplified. *Proc. Natl. Acad. Sci. USA* 105, 13486–13491.
- Stewart, J.B., Beckenbach, A.T., 2009. Characterization of mature mitochondrial transcripts in *Drosophila*, and the implications for the tRNA punctuation model in arthropods. *Gene* 445, 49–57.
- Suga, K., Welch, D.B.M., Tanaka, Y., Sakakura, Y., Hagiwara, A., 2008. Two circular chromosomes of unequal copy number make up the mitochondrial genome of the rotifer *Brachionus plicatilis*. *Mol. Biol. Evol.* 25, 1129–1137.
- Szitenberg, A., Rot, C., Ilan, M., Huchon, D., 2010. Diversity of sponge mitochondrial introns revealed by *cox1* sequences of Tetillidae. *BMC Evol. Biol.* 10, 288.
- Taanman, J.-W., 1999. The mitochondrial genome: structure, transcription, translation and replication. *Biochim. Biophys. Acta – Bioenerg.* 1410, 103–123.
- Temperley, R.J., Wydro, M., Lightowlers, R.N., Chrzanowska-Lightowlers, Z.M., 2010. Human mitochondrial mRNAs-like members of all families, similar but different. *Biochim. Biophys. Acta* 1797, 1081–1085.
- Thrash, J.C., Boyd, A., Huggett, M.J., Grote, J., Carini, P., Yoder, R.J., Robbertse, B., Spatafora, J.W., Rappé, M.S., Giovannoni, S.J., 2011. Phylogenomic evidence for a common ancestor of mitochondria and the SAR11 clade. *Sci. Rep.* 1, 13.
- Tiranti, V., Rocchi, M., DiDonato, S., Zeviani, M., 1993. Cloning of human and rat cDNAs encoding the mitochondrial single-stranded DNA-binding protein (SSB). *Gene* 126, 219–225.
- Torres, T.T., Dolezal, M., Schlötterer, C., Ottenwälder, B., 2009. Expression profiling of *Drosophila* mitochondrial genes via deep mRNA sequencing. *Nucl. Acids Res.* 37, 7509–7518.
- Vallès, Y., Halanych, K.M., Boore, J.L., 2008. Group II introns break new boundaries: presence in a bilaterian's genome. *PLoS ONE* 3, e1488.
- Viljakainen, L., Oliveira, D.C., Werren, J.H., Behura, S.K., 2010. Transfers of mitochondrial DNA to the nuclear genome in the wasp *Nasonia vitripennis*. *Insect Mol. Biol.* 19 Suppl. 1, 27–35.
- Voigt, O., Erpenbeck, D., Wörheide, G., 2008. A fragmented metazoan organellar genome: the two mitochondrial chromosomes of *Hydra magnipapillata*. *BMC Genom.* 9, 350.
- Walberg, M.W., Clayton, D.A., 1981. Sequence and properties of the human KB cell and mouse L cell D-loop regions of mitochondrial DNA. *Nucl. Acids Res.* 9, 5411–5421.
- Wang, X., Lavrov, D.V., 2008. Seventeen new complete mtDNA sequences reveal extensive mitochondrial genome evolution within the Demospongiae. *PLoS ONE* 3, e2723.
- Wang, X., Lavrov, D.V., 2011. Gene recruitment—a common mechanism in the evolution of transfer RNA gene families. *Gene* 475, 22–29.
- Wanrooij, P.H., Uhler, J.P., Simonsson, T., Falkenberg, M., Gustafsson, C.M., 2010. G-quadruplex structures in RNA stimulate mitochondrial transcription termination and primer formation. *Proc. Natl. Acad. Sci. USA* 107, 16072–16077.
- Watanabe, K., Yokobori, S.-i., 2011. tRNA modification and genetic code variations in animal mitochondria. *J. Nucl. Acids* 2011, 623095.
- Wei, S.-J., Shi, M., Chen, X.-X., Sharkey, M.J., van Achterberg, C., Ye, G.-Y., He, J.-H., 2010. New views on strand asymmetry in insect mitochondrial genomes. *PLoS ONE* 5, e12708.
- Wolstenholme, D.R., 1992. Animal mitochondrial DNA: structure and evolution. *Int. Rev. Cytol.* 141, 173–216.
- Wolstenholme, D.R., Okimoto, R., Macfarlane, J.L., 1994. Nucleotide correlations that suggest tertiary interactions in the TV-replacement loop-containing mitochondrial tRNAs of the nematodes, *Caenorhabditis elegans* and *Ascaris suum*. *Nucl. Acids Res.* 22, 4300–4306.
- Wong, T.W., Clayton, D.A., 1986. DNA primase of human mitochondria is associated with structural RNA that is essential for enzymatic activity. *Cell* 45, 817–825.
- Xu, W., Jameson, D., Tang, B., Higgs, P., 2006. The relationship between the rate of molecular evolution and the rate of genome rearrangement in animal mitochondrial genomes. *J. Mol. Evol.* 63, 375–392.
- Yamazaki, N., Ueshima, R., Terrett, J.A., Yokobori, S.I., Kaifu, M., Segawa, R., Kobayashi, T., Numachi, K.I., Ueda, T., Nishikawa, K., Watanabe, K., Thomas, R.H., 1997. Evolution of pulmonate gastropod mitochondrial genomes: comparisons of gene organizations of Euhadra, Cepaea and Albinaria and implications of unusual tRNA secondary structures. *Genetics* 145, 749–758.
- Yang, M.Y., Bowmaker, M., Reyes, A., Vergani, L., Angeli, P., Gringeri, E., Jacobs, H.T., Holt, I.J., 2002. Biased incorporation of ribonucleotides on the mitochondrial L-strand accounts for apparent strand-asymmetric DNA replication. *Cell* 111, 495–505.
- Zardoya, R., Meyer, A., 1996. Phylogenetic performance of mitochondrial protein-coding genes in resolving relationships among vertebrates. *Mol. Biol. Evol.* 13, 933–942.
- Zhang, D.-X., Hewitt, G.M., 1997. Insect mitochondrial control region: a review of its structure, evolution and usefulness in evolutionary studies. *Biochem. Syst. Ecol.* 25, 99–120.
- Zhong, M., Struck, T.H., Halanych, K.M., 2008. Phylogenetic information from three mitochondrial genomes of Terebelliformia (Annelida) worms and duplication of the methionine tRNA. *Gene* 416, 11–21.